arXiv:astro-ph/9810329v1 21 Oct 1998

A Model of the Double Magnetic Cycle of the Sun

E. E. Benevolenskaya
Pulkovo Astronomical Observatory, St. Petersburg, 196140, Russia
W.W.Hansen Experimental Physics Laboratory,
Stanford University, Stanford, CA 94305-4085

ABSTRACT

It has been argued that the solar magnetic cycle consists of two main periodic components: a low-frequency component (Hale's 22-year cycle) and a high-frequency component (quasi-biennial cycle). The existence of the double magnetic cycle on the Sun is confirmed using Stanford, Mount Wilson and Kitt Peak magnetograph data from 1976 to 1996 (solar cycles 21 and 22). In the frame of the Parker's dynamo theory a model of the double magnetic cycle is presented. This model is based on the idea of two dynamo sources separated in space. The first source of the dynamo action is located near the bottom of the convection zone, and the second operates near the top. The model is formulated in terms of two coupled systems of non-linear differential equations. It is demonstrated that in the case of weak interaction between the two dynamo sources the basic features of the double magnetic cycle such as existence of two component and observed temporal variations of high-frequency component can be reproduced.

Subject headings: Sun: magnetic field — Sun: activity — Suns interior

1. Introduction

The 22-yr cycle of solar activity is a magnetic cycle which consists of two 11-years sunspot cycles. The 22-yr cycles begin on even sunspot cycles according to the Zürich numbering (Gnevyshev and Ohl, 1948) and manifest themselves in reversal of polarity of sunspots (Hale's law). From one 11-yr cycle to another, the polarity of the preceding (p) and following (f) sunspots reverses. This reversal corresponds to the changing of the toroidal magnetic field (B_φ -component). During even-numbered cycles $B_\varphi > 0$ (the direction of the magnetic vector coincides with the direction of solar rotation) in the northern (N) hemisphere and $B_\varphi < 0$ in the southern (S) hemisphere. This relation is reversed during odd-numbered cycles. In parallel with this, the poloidal magnetic field, or background field (B_r -component), shows a 22-yr periodicity (Howard and LaBonte, 1981), and other studies (Stenflo, 1994) have suggested that the low-latitude and polar fields similarly show a 22-yr periodicity . Furthermore, these studies have hinted at a shorter term periodicity of about 2 yr (high-frequency component). In addition, during the periods of the three-fold polar magnetic field reversals which occur during some sunspot cycle maxima, the temporal separation of the zones of alternated polarity of the magnetic field on H_α charts is approximately equal to 1.5–2.5 years (Waldmeier, 1973, Benevolenskaya, 1991). Similar periodicities were found in variations of radio flux on 10.7cm (Belmont et al., 1966), flares and sunspot areas (Akioka et al., 1987). The high-frequency, 2-year, component is substantially weaker than the main 22-year cycle and its intensity varies with time. The two year component more clearly appeared in northern hemisphere in cycle 20 and in southern hemisphere in cycle 21; its value was smaller in cycle 22 (Benevolenskaya, 1996). However, the biennial cycle represents a challenge to the solar dynamo theories which usually explain only the main cycle. In this Letter, I present a new evidence of the two components of the solar cycle, and an attempt to explain the results in term of Parker's dynamo theory (Parker, 1979).

2. Analysis of magnetograph data

Using Kitt Peak magnetograph data we show how synoptic (longitude independent) magnetic field patterns evolve through cycles 21 and 22. Each of the synoptic maps is represented by the values of $B_\parallel$ as a function sine latitude and Carrington longitude. The Carrington coordinate system is a reference frame rotating rigidly at the rate which corresponds to the synodic rotation rate of sunspots at a latitude of about 16^o . The observed line-of-sight components ($B_\parallel$) averaging over all longitudes for each Carrington Rotation is represented in Figure 1b. The time step in the time set of values $B_\parallel$ corresponds to Carrington rotation period ($P = 27.2753$ days).

The relative sunspot number is shown in Figure 1a. Component B_r (Figure 1c), was calculated from observational data assuming that the true average field direction is radial and $B_\theta \cong 0$. The radial component will be easily found at all latitudes besides near the poles:

$$B_r = B_\parallel / \sin\theta, \qquad (1)$$

where θ is colatitude.

To separate the high-frequency component in the data, we apply a difference filter by computing $\Delta B_r/\Delta t$ for different time intervals Δt . This is a reasonable procedure if the poloidal magnetic field consists of two components and is represented as two dynamo waves:

$$B_r(t,\theta) = b_r(\theta)\sin\omega_c t + \frac{b_r(\theta)}{A}\sin(\omega_\delta t + \varphi). \qquad (2)$$

where $b_r(\theta)$ is the amplitude of the low-frequency radial component of the magnetic field; A is ratio between amplitudes of low-frequency and high-frequency components; ω_c is the frequency of the Hale's solar cycle; ω_δ is the frequency of biennial cycle; φ is a phase.

The expression for the $\Delta B_r/\Delta t$ can be written as

$$\frac{\Delta B_r(t,\theta)}{\Delta t} = b_r(\theta)\omega_c\cos\omega_c t + \frac{b_r(\theta)}{A}\omega_\delta\cos(\omega_\delta t + \varphi). \qquad (3)$$

When $\omega_\delta \cong 10\omega_c$, $A \cong 2$ as it took place in cycle 20, then the low-frequency term dominates in the expression for B_r (equation 2) and the high-frequency term prevails in the expression for $\Delta B_r/\Delta t$ (equation 3). Thus, presuming the existence of the double magnetic cycle, one finds that the low-frequency term prevails in the B_r -component, while the high - frequency term dominates in $\Delta B_r/\Delta t$.

Contour plots of $\Delta B_r/\Delta t$ as a function μ (or $\cos\theta$) and time, t, measured in Carrington rotation



Fig. 1.— (a) The values of the sunspot number for cycles 21 and 22 as function of time (in years). (b) The line-of-sight component of the solar magnetic field as function of Carrington rotation number. (c) The radial component of the solar magnetic field (blue lines correspond to isolines for +3, +10 G, red lines, for -3, -10 G). (d) The values of the $\Delta B_r/\Delta t$ for $\Delta t = 6$ solar Carrington rotations (blue lines correspond to isolines for +0.3, +3 G/rotation, red lines, for -0.3, -3 G/rotation).

are represented in Figure 1d for $\Delta t = 6P$. During the cycles 21 and 22 the zones of increasing and decreasing strength of the surface magnetic field appear in both the N and S hemispheres (see Figure 1d). The width of these zones is approximately 2 years. This result coincides with investigations done earlier based on Wilcox and Mount Wilson synoptic maps. Corresponding power spectra for $\Delta B_r/\Delta t$ show that the component with period $T = 1.5 - 2.5$ years is dominant in both hemispheres (Benevolenskaya, 1995, Benevolenskaya, 1996). These results are the basis of our suggestion of a two component of solar cycle.

3. Model

The possible explanation of the double magnetic cycle is that the magnetic fields are generated by Parker's dynamo acting in convective zone. The low-frequency component is generated at the base of the convective zone due to large scale radial shear $\partial\Omega/\partial r$; Ω is the angular velocity. The high-frequency component may be generated in subsurface regions due to latitudinal shear $\partial\Omega/\partial\theta$ or due to radial shear. The recent investigations of solar interior rotation show a significant radial gradient of angular velocity exists in subsurface of the convective zone together with the latitudinal gradient of the angular velocity (Schou, et al., 1998). For simplicity we use only latitudinal shear for generation of the high-frequency component. Cartesian coordinates are employed, with x denoting the radial, y the azimuthal and z the latitudinal coordinate. We consider axisymmetrical solutions ($\dfrac{\partial}{\partial\varphi} = 0$).

At the base of the convection zone turbulence is suppressed by strong magnetic field (Parker, 1993) and, therefore, diffusivity in the first layer η_1 will could be less then η_2 in the second layer. The axisymmetrical mean magnetic field is decomposed into toroidal and poloidal parts. The following set of equations describe evolution of the magnetic field in thin layers at the levels x_1 and x_2:

$$\begin{cases} \dfrac{\partial A_{y1}}{\partial t} &= (\alpha_1 + C_1)B_{y1} + \eta_1 \Delta A_{y1} \\ \dfrac{\partial B_{y1}}{\partial t} &= -\dfrac{\partial A_{y1}}{\partial z} G_x + \eta_1 \Delta B_{y1} \\ \dfrac{\partial C_1}{\partial t} &= -\nu C_1 + p A_{y1} B_{y1} \end{cases} \tag{4}$$

$$\begin{cases} \dfrac{\partial A_{y2}}{\partial t} &= (\alpha_2 + C_2)B_{y2} + \eta_2 \Delta A_{y2} \\ \dfrac{\partial B_{y2}}{\partial t} &= \dfrac{\partial A_{y2}}{\partial x} G_z + \eta_2 \Delta B_{y2} \\ \dfrac{\partial C_2}{\partial t} &= -\nu C_2 + p A_{y2} B_{y2} \end{cases} \tag{5}$$

where B_y is the toroidal magnetic field and A_y is the azimuthal vector potential which gives poloidal field, η is diffusivity and α is a kinetic helicity, C is the variable part of kinetic helicity. The first and the second equations in the both systems (4) and (5) are a generation of Parker's equations for large-scale radial shear $G_x = \dfrac{\partial V_y}{\partial x}$ and large-scale latitudinal shear $G_z = \dfrac{\partial V_y}{\partial z}$ correspondingly. We have used the third equation in the both systems $\dfrac{\partial C}{\partial t} = -\nu C + pAB$ according to Kleeorin and Ruzmaikin (1982) in a simple form, for feedback of the magnetic field on the helicity. In these equations $\nu > 0$ and $p < 0$ are parameters of the feedback. These two non-linear differential equation systems describe the evolution of two independent sources of the magnetic field. In the frame of this model it is difficult to explain observed variations of the high-frequency component. Therefore, we assume that the erupted low-frequency magnetic field can influence the physical conditions in the region where high-frequency component operates, through modifying the helicity there. This allows the high-frequency component to vary with time. In this case, the equation for C_2 becomes $\dfrac{\partial C_2}{\partial t} = -\nu C_2 + p(A_{y2} + aA_{y1})(B_{y2} + aB_{y1})$, where the parameter a captures the feedback of the low-frequency component of the magnetic field on the helicity in regions where the high-frequency component is found. Following Weiss et al. (1984) the solutions of these equations were found as $A_1 = a(t)e^{ikx}, B_1 = b(t)e^{ikx}$ and $C_1 = c_o(t) + c(t)e^{2ikx}$, Im $c_o = 0$ for x_1 level and the same expressions for x_2 level where x is replaced by z. A, B, C are complex functions as $A = a_1 + ia_2$, $B = b_1 + ib_2$.

It is convenient to present the system in dimensionless units. Since there will be an underlying periodicity in all calculations, we use those periods as relative units of time. In relative units $t \longrightarrow t^{'} = \eta_2 k^2 t \alpha_o$, $\alpha \longrightarrow \alpha^{'} = \alpha/\eta_2 k^2 \alpha_o$, $\sigma = \eta 1/\eta 2$, $C \longrightarrow C^{'} = C/\eta_2 k^2 \alpha_o, G_z \longrightarrow G_z^{'} = G_z/\eta_2 k \alpha_o$ and $G_x \longrightarrow G_x^{'} = G_x/\eta_2 k \alpha_o$, where α_o is some mean value of the kinetic helicity. Therefore, G_x and G_z are represented as dynamo numbers in our model. Therefore, set of the partial differential equations reduced to the following equations were then solved numerically.

$$\begin{cases} \dot{A}_1 &= (\alpha_1 + C_1)B_1 - \sigma A_1 + \frac{1}{2} B_1^* \, C_1 \\ \dot{B}_1 &= -iA_1^* \, G_x - \sigma B_1 \\ \dot{C}_1 &= -\nu C_1 + pA_1 B_1 \end{cases} \tag{6}$$

$$\begin{cases} \dot{A}_2 &= (\alpha_2 + C_2)B_2 - A_2 + \frac{1}{2} B_2^* \, C_2 \\ \dot{B}_2 &= iA_2 G_z - B_2 \\ \dot{C}_2 &= -\nu C_2 + p(A_2 + aA_1)(B_2 + aB_1) \end{cases} \tag{7}$$

We have investigated solutions of these two non-linear systems as functions of the parameter a. Results for three principal cases, $a = 0$, $0 < a < 2$ and $a = 2$ are represented in Figure 3, where $\alpha_1 = \alpha_2 = 1$, $\nu = 0.5$, $p = -1$, $\sigma = 0.1$, $G_x = 0.3$, $G_z = 20$, which are in the reasonable range for the Sun. If $a = 0$ we have two independent sources of the magnetic field which together generate low-frequency and high-frequency signatures (Fig. 2 a, b). For a weak interaction between these sources ($0 < a < 0.2$) we still get a low and high frequencies response. (Fig. 2 c, d). Moreover, the high- frequency component becomes more regular, and consequently is decreased when a increases. With further increase of the feedback of the low-frequency magnetic field on the helicity near the top surface the high-frequency component disappears and only a low-frequency mode regime is established (Fig. 2 e, f). Therefore, in the case of weak interaction between two sources of magnetic field it is possible to obtain a stable double magnetic cycle.

4. Conclusions

Our model simulates the double magnetic cycle and temporal variations of the biennial cycles from one 11-year cycle to another. According to the magnetograph data (Benevolenskaya, 1996) the 2-year component more clearly appeared in the northern hemisphere in cycle 20 and in the southern hemisphere in cycle 21. In cycle 22 it was present in both hemispheres but was of lower amplitude.

In our model, a reduced high-frequency component occurs when the erupted magnetic field of the main (Hale's) cycle imposes the helicity in the regions of generation of the high-frequency component. The high-frequency component is more pronounced when the effect of erupting magnetic fields of the main cycle on helicity in this region is small. Thus, this simple model provides a qualitative explanation of the double magnetic cycle. The next step to a quantitative model is in development of a 2D model in spherical geometry, since this simple model gives only a qualitative picture of the two-components of the solar magnetic cycle.

Acknowledgments.

I am grateful to Dr. J.W.Harvey for providing us the Kitt Peak magnetograph data and to Drs. P.A.Gilman, J.T.Hoeksema, A.G.Kosovichev and P.H.Scherrer for useful discussions and Russian Federal Programme “Astronomy”, grant 1.5.3.4.

REFERENCES

Akioka, M., Kubota, J., Suzuki, M. et al. 1987, Sol. Phys., **112**, 313

Belmont, A.D., Darff, D.C., Ultad, M.S. 1966, *J. Atmos.Sci.*, **23**, 314

Benevolenskaya, E.E. 1991, in *The Sun and Cool Stars: Activity, Magnetism, Dynamos*, Tuominen, I., Moss, D. and Rüdiger, G. (eds.), Springer-Verlag, 234

Benevolenskaya, E.E. 1995, Sol. Phys., **161**, 1

Benevolenskaya, E.E. 1996, Sol. Phys., **167**, 47

Gnevyshev, M.N., and Ohl, A.I. 1948, AZh, **25**, 18

Howard, R., and LaBonte, B.J. 1981, Sol. Phys., **74**, 131

Kleeorin, N.I., and Ruzmaikin, A.A. 1982 *Magnitnaya Gidrodinamika*, No 2, 17

Parker, E.N. 1979, *Cosmic Magnetic Fields*, Oxford University Press

Fig. 2.— (a) Real part of $A = A_1 + A_2$ as function time for independent systems ($a = 0$). (b) Power Spectrum for case (a) in relative units. (c) Real part of $A = A_1 + A_2$ as function time for $a = 0.17$. (d) Power Spectrum for case (c) in relative units. (e) Real part of $A = A_1 + A_2$ as function time for $a = 0.2$. (f) Power Spectrum for case (e) in relative units.

Parker, E.N. 1993, ApJ, **408**, 707

Schou, J., et al., 1998, ApJ, in press

Stenflo, J.O. 1994, in *Solar Surface Magnetism*, Rutten, R.J. and Schrijver, C.J. (eds.), NATO Advanced Research Workshop, Kluwer

Waldmeier, M. 1973, Sol. Phys., **28**, 389

Weiss, N.O., Cattaneo, F., Jones, C.A. 1984, *Geophys. Astrophys. Fluid. Dyn.*, **30**, 305

This 2-column preprint was prepared with the AAS LATEX macros v4.0.